

North Yorkshire Joint Strategic Needs Assessment 2019 Airedale, Wharfedale and Craven CCG Profile

Introduction

This profile provides an overview of population health needs in Airedale, Wharfedale and Craven CCG (AWC CCG). Greater detail on particular topics can be found in our Joint Strategic Needs Assessment (JSNA) resource at www.datanorthyorkshire.org which is broken down by district. This document is structured into five parts: population, deprivation, disease prevalence, hospital admissions and mortality. It identifies the major themes which affect health in AWC CCG and presents the latest available data, so the dates vary between indicators.

Summary

- **Life expectancy is not significantly different from England.** For 2011-2015, female life expectancy in AWC CCG is 83.3 years (England: 83.1), and male life expectancy is more than three years lower than for females at 79.8 years (England: 79.4) [1].
- **There is a high proportion of older people.** In 2017, 21.2% of the population was aged 65 and over (33,600), higher than national average (17.3%). Furthermore over 4,800 (3.0%) were age 85+, compared with 2.3% in England. [2]
- **Some children grow up in relative poverty.** In 2015, there were 14.9% of children aged 0-15 years living in low income families, compared with 19.9% in England [1].
- **There are areas of deprivation.** Within the CCG, 19 Lower Super Output Areas (LSOAs) out of a total of 97 are amongst the 20% most deprived in England, and 11 out of these 19 LSOAs are amongst the 10% most deprived in England. These 11 LSOAs are all in the Keighley Central, Keighley West and Keighley East wards in Bradford [3].
- **Many people have longstanding health problems.** The census in 2011 showed 27,500 people living with long-term health problem or disability (17.4% compared to 17.6% in England) [1].
- **The highest reported rates of ill health are from:** hypertension (15.3%); obesity (11.6%); depression (10.7%); diabetes (7.3%); and asthma (7.2%) [4].
- **Hospital admissions vary according to admissions route.** Non-elective admissions are most frequently due to respiratory problems (15.7%); injury, poisoning and certain other consequences of external causes (11.9%); and circulatory diseases (11%). Elective admissions are most common for neoplasms (19.6%); factors influencing health status (13.6%); and digestive disorders (13%) [5].


Population

There are 16 general practices in AWC CCG area with 159,000 [registered patients](#) (December 2018) [6]. In contrast, the ONS mid-year resident population estimate for 2017 gave a CCG-wide population of 159,800 [7]. The GP registered population in AWC CCG is 0.5% lower than the resident population, whereas in England, the registered population is 7% higher than the resident population.

The resident population is forecast to rise to 164,400 by 2025 (2.3% increase since 2018) and 169,400 by 2040 (5.4% increase since 2018) [8]. In England, the corresponding increases are 4% by 2025 and 10.3% by 2040. Local population growth is forecast to be lower than that seen nationally.

There is a high proportion of people aged over 65 (21.2%) in the AWC CCG compared with England (17.3%). The proportion of people aged 5-14 (11.9%) is slightly higher than England (11.6%). The following age profile shows a lower proportion of the population in age groups 0-4 and 20-39 years, and a higher proportion in age groups 50-95+, compared with both England and the Yorkshire & Humber region.

Age Profile – GP registered population by sex and five-year age band 2017




Source: National General Practice Profiles, PHE

Deprivation

In 2015, there were 14.9% of children aged 0-15 years living in low income families, compared with 19.9% in England [1]. The 2015 [Index of Multiple Deprivation](#) (IMD) identifies 19 Lower Super Output Areas (LSOAs) out of a total of 97 across the CCG which are amongst the 20% most deprived in England, and 11 out of these 19 LSOAs are amongst the 10% most deprived in England. These 11 LSOAs are all in the Keighley Central, Keighley West and Keighley East areas in Bradford. Two LSOAs in Skipton South, Craven are amongst the 19 LSOAs (20% most deprived in England) [3]. A list of these 19 LSOAs can be found in Appendix 1.

Deprivation scores, using IMD-2015, have been estimated for general practices. They show five practices in AWC CCG have populations experiencing higher levels of deprivation than England, although they are not within North Yorkshire's boundary.



Source: National General Practice Profiles, PHE

* Practices within North Yorkshire.


IG Medical is based in Ilkley (Bradford), with a branch in Grassington (North Yorkshire)

Lifestyle and behaviour

The lifestyle choices that people make and behaviours they follow in their lifetime can all have an impact on both their current and future health. Lifestyle diseases are defined as diseases linked with the way people live their life. This is commonly caused by alcohol, drug and smoking abuse as well as lack of physical activity and unhealthy eating.

Smoking


AWC CCG has the second highest estimated rate of smoking in North Yorkshire, after Scarborough & Ryedale CCG, in 2017/18. Although the rate is high when compared to other CCGs in North Yorkshire, AWC CCG has a lower rate of smoking prevalence compared to England. Five practices have rates which are significantly higher than England and nine practices have significantly lower rates than CCG and England averages.



Source: National General Practice Profiles, PHE

Adult obesity


There is a higher rate of adult obesity in AWC CCG compared with England, with around 14,700 people on the register. Seven out of the 16 practices have rates which are significantly higher than both the CCG average and England and seven practices have rates significantly lower than the CCG average. Four practices have rates significantly lower than England.



Source: NHS Digital

Disease Prevalence

In AWC CCG, hypertension, obesity and depression are the most common health problems, followed by diabetes and asthma. The prevalence for most diseases and risk factors is higher in AWC CCG than for England.




Source: NHS Digital

Disease prevalence by general practice

The following charts use the NHS Quality and Outcomes Framework prevalence data for 2017/18. These are expressed as crude percentages, without taking account of variation in the populations between general practices. Differences such as the proportion of elderly patients, ethnicity and levels of deprivation may affect crude prevalence rates. The charts are presented in order of recorded prevalence, from highest to lowest. Practices within North Yorkshire are indicated by an asterisk (*).

Hypertension


In AWC CCG, there are around 24,400 people with known hypertension and prevalence is higher than England. Twelve general practices have rates significantly higher than England, whilst four have significantly lower rates.



Source: NHS Digital

Depression


There are more than 13,500 people with a record of depression in AWC CCG, which is a slightly higher rate than seen in England. Eight practices have rates which are significantly higher than England, while six practices have significantly lower rates than both England and the CCG average.



Source: NHS Digital

Asthma


In AWC CCG, asthma prevalence is higher than England. There are around 11,500 people on asthma registers. Nine out of 16 practices have asthma prevalence rates which are significantly higher than England. The other seven practices have a recorded prevalence which is similar to England.



Source: NHS Digital

Coronary heart disease

Coronary heart disease (CHD) prevalence is higher in AWC CCG compared with England and there are around 6,400 people with diagnosed CHD. Most general practices (13 of the 16) have prevalence rates significantly higher than England. Silsden & Steeton Medical Practice has a prevalence rate 50% higher than England.



Source: NHS Digital

Consideration can be given to variation which may be due to modifiable risk factors within the population, differences in record keeping, variation in health care and access to services. [NHS RightCare](#) produces a range of intelligence products to help local health economies identify and address health inequality.

Furthermore, the NHS Health Check is a health check-up for adults in England aged 40-74, designed to spot early signs of stroke, kidney disease, heart disease, type 2 diabetes or dementia. As individuals age they have a higher risk of developing one of these conditions and an NHS Health Check helps find ways to lower this risk. The NHS Health Check report for North Yorkshire highlights performance of health checks across North Yorkshire and can be found on [Data North Yorkshire](#).


Hospital admissions

In 2016/17, there were just over 63,000 hospital admissions for AWC CCG registered patients, of which 41,553 (65.5%) were elective admissions and 21,909 (34.5%) were non-elective admissions. In total, there were 126 providers, with Airedale NHS Foundation Trust being the main provider.

Hospital admissions by provider, AWC CCG, 2016/17			
Provider	Proportion of elective admissions	Proportion of non-elective admissions	Proportion of all admissions
Airedale NHS Foundation Trust	74.8%	86.8%	78.9%
Bradford Teaching Hospitals NHS Foundation Trust	7.9%	5.8%	7.1%
Leeds Teaching Hospitals NHS Trust	7.0%	4.6%	6.2%
Ramsay Healthcare UK Operations Ltd	6.5%		4.3%
Harrogate & District NHS Foundation Trust	1.1%		
Remaining providers	2.8%	2.8%	3.5%

Source: Public Health England SHAPE atlas


The main reasons for non-elective admissions are shown below for causes which contributed towards more than 5% of non-elective admissions. Respiratory diseases are the most common reason for non-elective admission followed by injuries & poisoning and circulatory diseases.



Source: Public Health England SHAPE atlas

Within *chapter XIX: Injury, poisoning and certain other consequences of external causes*, the main reasons for admission are: poisoning by non-opioid drugs; fracture of femur; fracture of lower leg; and superficial head injury. This suggests drug overdose (accidental or otherwise) and falls may contribute importantly to local emergency admissions.

The main reasons for elective admission are similarly shown for causes which contributed towards more than 5% of elective admissions. Neoplasms represent the highest proportion of elective admissions, followed by factors influencing health status and digestive diseases.




Source: Public Health England SHAPE atlas

For *chapter XXI: Factors influencing health status*, the leading reasons for admission are: supervision of normal pregnancy; liveborn infants according to place of birth; special screening for neoplasms; and follow-up examination after treatment for malignant neoplasm.


Under 18 hospital admissions

AWC CCG has a slightly higher rate of admissions due to injury for those under compared to England. Over half of practices in AWC CCG have higher rates than the England and CCG average.



Source: National General Practice Profiles, PHE

AWC CCG has the highest rate of emergency hospital admissions for all causes under 18 when compared to other CCGs in North Yorkshire. The rate is also higher than the England average. However, AWC CCG has a rate of A&E attendance under 18 which is lower than the England average.




Source: National General Practice Profiles, PHE

Public Health England produces a summary health profile for AWC CCG (Appendix 2). This compares more than 50 indicators with national data and highlights those which are significantly different from England. This can be used to help inform topics which might be considered for focused improvement work. In particular, it highlights the following as being significantly worse than England:

- Provision of 1 hour or more unpaid care per week (%)
- Emergency admissions in under 5s (Crude rate per 1,000)
- Emergency hospital admissions for CHD (SAR)
- Hospital stays for alcohol related harm (SAR)
- Elective hospital admissions for hip replacement (SAR)

Mortality


The chart below shows the directly standardised rate (DSR) of potential years of life lost (PYLL) per 100,000 registered patients and the number of observed deaths by conditions. The condition with the highest DSR is Neoplasm (730.8), while the highest number of observed deaths is from ischaemic heart diseases (154).



Source: HSCIC

Place of death

Within AWC CCG, 35.2% of deaths occurred in hospital, 29.4% in care homes, 22.9% at home, 9.0% in hospices and 3.5% elsewhere. Compared with England, AWC CCG has fewer people dying in hospital and at home but less people dying in care homes and hospices [9].



Source: Public Health England

Additional mortality data available in the [JSNA 2018 District Profiles](#).

References

1. Public Health England. [Local Health](#)
2. Public Health England. [National General Practice Profiles](#)
3. [Data.gov.uk](#)
4. NHS Digital. QOF 2017/18
5. [SHAPE](#) (registration required)
6. NHS Digital. [CCG outcomes tool](#)
7. ONS. [Clinical commissioning group population estimates](#)
8. ONS. [Population projections - clinical commissioning groups](#)
9. Public Health England. [End of Life Care Profiles](#)

Contributors:

Judith Yung, Public Health Intelligence Analyst

Emel Perry, Public Health Intelligence Analyst

Wendy Rice, Public Health Intelligence Analyst

Sharon Draper, Data Officer

Katie Wilkinson, Data Officer

Leon Green, Senior Public Health Intelligence Specialist

Contact:

nypublichealth@northyorks.gov.uk

January 2019

Appendix 1

LSOA	Ward	District	Index of Multiple Deprivation (IMD) National Rank (where 1 is most deprived)	Index of Multiple Deprivation (IMD) Decile (where 1 is most deprived 10% of LSOAs)
Bradford 011A	Keighley Central	Bradford	60	1
Bradford 008G	Keighley Central	Bradford	284	1
Bradford 008B	Keighley Central	Bradford	691	1
Bradford 009D	Keighley Central	Bradford	1,214	1
Bradford 009C	Keighley West	Bradford	1,541	1
Bradford 008A	Keighley East	Bradford	1,772	1
Bradford 009B	Keighley West	Bradford	2,088	1
Bradford 011B	Keighley East	Bradford	2,169	1
Bradford 008C	Keighley Central	Bradford	2,777	1
Bradford 007A	Keighley Central	Bradford	2,832	1
Bradford 009A	Keighley Central	Bradford	2,917	1
Bradford 012C	Keighley West	Bradford	3,298	2
Bradford 014A	Keighley West	Bradford	3,552	2
Bradford 007C	Keighley Central	Bradford	3,893	2
Bradford 009E	Keighley West	Bradford	3,943	2
Bradford 011D	Keighley East	Bradford	5,255	2
Craven 006B	Skipton South	Craven	5,801	2
Craven 005D	Skipton South	Craven	6,114	2
Bradford 012A	Keighley West	Bradford	6,187	2

Appendix 2

AWC CCG health profile summary

Selection: E3800001 - NHS Airedale, Wharfedale and Craven CCG

Indicators	Selection value	England value	England worst	Summary chart	England best
Low Birth Weight of term babies (%)	3.1	2.8	5		1.7
Child Development at age 5 (%)	60.4	60.4	42.2		74.3
GCSE Achievement (5A*-C inc. Eng & Maths) (%)	56.3	56.6	40.2		75.4
Unemployment (%)	1.4	1.8	4.8		0.5
Long Term Unemployment (Rate/1,000 working age population)	2.8	3.7	14.4		0.5
General Health - bad or very bad (%)	4.8	5.5	9.5		2.8
General Health - very bad (%)	1	1.2	2.2		0.6
Limiting long term illness or disability (%)	17.4	17.6	25.6		11.2
Overcrowding (%)	5.1	8.7	34.9		2.7
Provision of 1 hour or more unpaid care per week (%)	10.8	10.2	13		6.5
Provision of 50 hours or more unpaid care per week (%)	2.2	2.4	4		1.3
Pensioners living alone (%)	31.3	31.5	45.2		25.7
Deliveries to teenage mothers (%)	1.1	1.1	2.3		0.2
Emergency admissions in under 5s (Crude rate per 1000)	237.2	149.2	269.8		65.3
A&E attendances in under 5s (Crude rate per 1000)	400.9	551.6	1719.5		221
Admissions for injuries in under 5s (Crude rate per 10,000)	138.3	138.8	280.3		77.7
Admissions for injuries in under 15s (Crude rate per 10,000)	108.4	110.1	183.9		65.2
Admissions for injuries in 15 - 24 year olds (Crude rate per 10,000)	144.6	137	238.6		53.9
Occasional smoker (modelled prevalence, age 15) (%)	4.1	4	5.3		1.2
Regular smoker (modelled prevalence, age 15) (%)	8.2	8.7	12.7		3.2
Obese adults (%)	23	24.1	30.9		14.5
Binge drinking adults (%)	22.3	20	34.5		7.5
Healthy eating adults (%)	28.4	28.7	19.4		46.5
Obese Children (Reception Year) (%)	7.4	9.3	13.8		5.3
Children with excess weight (Reception Year) (%)	20.2	22.2	27.2		14.6
Obese Children (Year 6) (%)	16.9	19.3	27.1		9.8
Children with excess weight (Year 6) (%)	30	33.6	42.9		21.7
Emergency hospital admissions for all causes (SAR)	98.7	100	152.5		68.2
Emergency hospital admissions for CHD (SAR)	115.9	100	270.9		59.4
Emergency hospital admissions for stroke (SAR)	93.7	100	160.1		76.8
Emergency hospital admissions for Myocardial Infarction (heart attack) (SAR)	103.8	100	280.2		53.8
Emergency hospital admissions for Chronic Obstructive Pulmonary Disease (COPD) (SAR)	86.4	100	233.4		43.2
Incidence of all cancer (SIR)	101.9	100	118.7		84.5
Incidence of breast cancer (SIR)	96.5	100	119.3		76.4
Incidence of colorectal cancer (SIR)	99.5	100	120.3		76.6
Incidence of lung cancer (SIR)	93	100	191.6		57
Incidence of prostate cancer (SIR)	101.2	100	147.4		64.3
Hospital stays for self harm (SAR)	98.9	100	308.4		28.7
Hospital stays for alcohol related harm (SAR)	108.6	100	182.5		57.7
Emergency hospital admissions for hip fracture in 65+ (SAR)	94.4	100	123.3		72.6
Elective hospital admissions for hip replacement (SAR)	109.7	100	140.9		32.7
Elective hospital admissions for knee replacement (SAR)	98.7	100	149.4		36.4
Life expectancy at birth for males, 2011- 2015 (years)	79.8	79.4	73.9		82.4
Life expectancy at birth for females, 2011- 2015 (years)	83.3	83.1	78.8		86
Deaths from all causes, all ages (SMR)	98.3	100	147.9		75.5
Deaths from all causes, under 65 years (SMR)	90	100	179.1		69.3
Deaths from all causes, under 75 years (SMR)	91.9	100	177		72.7
Deaths from all cancer, all ages (SMR)	98.5	100	127.9		78.3
Deaths from all cancer, under 75 years (SMR)	94.9	100	136.5		76.4
Deaths from circulatory disease, all ages (SMR)	97.6	100	153.5		73.1
Deaths from circulatory disease, under 75 years (SMR)	88.3	100	230.1		61.5
Deaths from coronary heart disease, all ages (SMR)	96.1	100	176.8		66.3
Deaths from coronary heart disease, under 75 years (SMR)	89.9	100	245.4		50.8
Deaths from stroke, all ages (SMR)	106.6	100	174.5		67.3
Deaths from respiratory diseases, all ages (SMR)	88.9	100	177.9		70.5

significantly worse significantly better not significantly different from average

Appendix 3

AWC CCG Outcomes Framework

● In IQ Range
 ● In best quartile
 ● CCG
 ◆ Cluster mean
 | England mean

Indicator Name	Value	Spine chart
CCG Outcomes Indicator Set- domain 1		
1.1 Potential years of life lost (PYLL) from causes considered amenable to healthcare - Female (2014)	2,288 ● ↑	1055 3204
1.1 Potential years of life lost (PYLL) from causes considered amenable to healthcare - Male (2014)	2,302 ● ↑	1325 3902
1.2 Under 75 mortality rates from cardiovascular disease (2016)	63.1 ● ↑	39.7 151.2
1.3 Completion of cardiac rehabilitation following an admission for coronary heart disease (2013/14)	No Data	0 75.4
1.4 Myocardial infarction, stroke and stage 5 chronic kidney disease in people with diabetes (2015/16)	86.2 ● ↑	52.3 280.2
1.5 Mortality within 30 days of hospital admission for stroke (2016/17)	1.48 ● ↑	0.29 1.62
1.6 Under 75 mortality rates from respiratory disease (2016)	32.9 ● ↑	15.1 88.6
1.7 Under 75 mortality rates from liver disease (2016)	16.3 ● ↑	7.1 41.4
1.8 Emergency admissions for alcohol related liver disease (2017 - 2017 (Jan - Dec))	45.5 ● ↑	7.2 79.3
1.9 Under 75 mortality rates from cancer (2016)	127 ● ↑	77 307
1.10 One-year survival from all cancers (Diagnosed 2015)	73.1 ● ↑	67 77.4
1.11 One-year survival from breast, lung and colorectal cancers (Diagnosed 2011)	69.1 ● ↑	62.1 76.2
1.12 People with Serious Mental Illness (SMI) who have received the complete list of physical checks (2014/15)	27.7 ● ↓	17.5 52.4
1.14 Maternal smoking at delivery (2017/18 Q3)	11.73 ● ↑	1.62 27.85
1.15 Breast feeding prevalence at 6 - 8 weeks (2015/16 Q1)	No Data	0 82.5
1.17 Record of stage of cancer at diagnosis (2016)	84.2 ● ↑	66.1 86.8
1.18 Percentage of cancers detected at stage 1 and 2 (2016)	52.9 ● ↑	39.4 60.4
1.19 Record of lung cancer stage at decision to treat (2016)	96.9 ● ↑	74.5 99.2
1.20 Mortality from breast cancer in females (2014 - 2016)	36.1 ● ↓	22.1 48
1.21 All-cause mortality - 12 months following a first emergency admission to hospital for heart failure in people aged 16 and over (April 2013 to March 2016)	94.0 ● ↓	75.6 119.7
1.22 Hip fracture: incidence (2017 - 2017 (Jan - Dec))	391 ● ↓	64 626
1.23 Smoking rates in people with serious mental illness (SMI) (2014/15)	39.7 ● ↓	27.2 55
1.24 Referrals to cardiac rehabilitation within 5 days of an admission for coronary heart disease (2014/15)	No Data	0 41.9
1.25 Neonatal mortality and stillbirths (2016)	4.70 ● ↓	2 13.3
1.26 Low birth weight full-term babies (2016)	2.20 ● ↓	1.3 5.2
CCG Outcomes Indicator Set- domain 2		
2.1 Health-related quality of life for people with long-term conditions (2016/17)	0.76 ● ↑	0.64 0.82
2.2 Proportion of people who are feeling supported to manage their condition (2016/17)	70.7 ● ↑	52.1 74.2
2.3 The percentage of people with Chronic Obstructive Pulmonary Disease (COPD) and Medical Research Council (MRC) Dyspnoea Scale >=3, identified on GP systems, referred to a pulmonary rehabilitation (2014/15)	35.5 ● ↑	3.8 68.5
2.4 Percentage of people with diabetes who have received nine care processes (2016/17)	45.0 ● ↓	17.7 86.9
2.5 People with diabetes diagnosed less than a year referred to structured education (2014/15)	68.8 ● ↑	41.7 93.2
2.6 Unplanned hospitalisation for chronic ambulatory care sensitive conditions (2017 - 2017 (Jan - Dec))	835 ● ↑	177 1390
2.7 Unplanned hospitalisation for asthma, diabetes and epilepsy in under 19s (2017 - 2017 (Jan - Dec))	390 ● ↑	40 647
2.8 Complications associated with diabetes (2015/16)	115.1 ● ↑	62.3 278.8
2.9 Access to community mental health services by people from Black and Minority Ethnic (BME) groups (2014/15)	1,785 ● ↑	658 5283
2.10 Access to psychological therapies services by people from Black and Minority Ethnic (BME) groups (2015/16)	1,198 ● ↑	295 13013
2.11a Percentage of referrals to Improving Access to Psychological Therapies (IAPT) services which indicated a reliable recovery following completion of treatment (2015 - 2015 (Jan - Dec))	40.7 ● ↑	18.7 60.5
2.11b Percentage of referrals to Improving Access to Psychological Therapies (IAPT) services which indicated a reliable improvement following completion of treatment (2015 - 2015 (Jan - Dec))	59.4 ● ↑	33.6 79.8
2.11c Percentage of referrals to Improving Access to Psychological Therapies (IAPT) services which indicate a reliable deterioration following completion of treatment (2015 - 2015 (Jan - Dec))	8.10 ● ↑	3.4 11.3
2.15 Health-related quality of life for carers, aged 18 and above (2016/17)	0.78 ● ↑	0.73 0.85
2.16 Health-related quality of life for people with a long-term mental health condition (2016/17)	0.49 ● ↑	0.34 0.68

CCG Outcomes Indicator Set - domain 3				
3.1 Emergency admissions for acute conditions that should not usually require hospital admission (2017 - 2017 (Jan - Dec))	1,590 ● ↓		225	2115
3.2 Emergency readmissions within 30 days of discharge from hospital (2011/12)	11.0 ● ↓		8.9	14.5
3.3 Elective Hip replacement (Primary) procedures - patient reported outcomes measures (PROMS) (2015/16)	0.46 ● ↑		0.35	0.52
3.3 Elective knee replacement (Primary) procedures - patient reported outcomes measures (PROMS) (2015/16)	0.34 ● ↑		0.19	0.39
3.3 Elective groin hernia procedures - patient reported outcomes measures (PROMS) (2015/16)	0.09 ● ↓		0.04	0.15
3.3 Elective varicose veins procedures - patient reported outcomes measures (PROMS) (2015/16)	No Data		0	0.15
3.4 Emergency admissions for children with lower respiratory tract infections (2017 - 2017 (Jan - Dec))	676 ● ↓		39	838
3.5 People who have had a stroke who are admitted to an acute stroke unit within 4 hours of arrival to hospital (2016/17)	52.8 ● ↓		17.6	85.1
3.6 People who have had an acute stroke who receive thrombolysis (2016/17)	7.60 ● ↓		0	27.6
3.7 People with stroke who are discharged from hospital with a joint health and social care plan (2016/17)	98.6 ● ↑		34.3	100
3.8 People who have a follow-up assessment between 4 and 8 months after initial admission for stroke (2016/17)	73.30 ● ↑		0	96.1
3.9 People who have had an acute stroke who spend 90% or more of their stay on a stroke unit (2016/17)	78.9 ● ↓		57.7	97.9
3.10.i Hip fracture: proportion of patients recovering to their previous levels of mobility/walking ability at 30 days (2015)	No Data		0	88.9
3.10.ii Hip fracture: proportion of patients recovering to their previous levels of mobility/walking ability at 120 days (2016)	No Data		41.1	92.9
3.11 Hip fracture: collaborative orthogeriatric care (2016)	90.7 ● ↑		55.1	100
3.12 Hip fracture: timely surgery (2016)	71.4 ● ↓		40.1	90.6
3.13 Hip fracture: multifactorial falls risk assessment (2016)	94.4 ● ↓		73.9	100
3.14 Alcohol-specific hospital admissions (2017 - 2017 (Jan - Dec))	102.6 ● ↓		33.9	322.6
3.15 Emergency alcohol-specific readmission to any hospital within 30 days of discharge following an alcohol-specific admission (2015 - 2017 (Jan - Dec))	83.1 ● ↓		41.9	198.2
3.16 Unplanned readmissions to mental health services within 30 days of a mental health inpatient discharge in people aged 17 and over (2014/15)	82.1 ● ↓		20.9	317.5
3.17 Percentage of adults in contact with secondary mental health services in employment (2016 - 2017 (Dec - Dec))	7.00 ● ↑		0	24
3.18 Hip fracture: care process composite indicator (2016)	54.8 ● ↓		25.5	87.9

CCG Outcomes Indicator Set - domain 4				
4.1 Patient experience of GP out-of-hours services (2014/15)	66.6 ● ↓		49	85.3
4.2 Patient experience of hospital care (2015/16)	77.8 ● ↑		68.3	83.5
4.5 Responsiveness to Inpatients personal needs (2015/16)	71.2 ● ↑		60.1	78

CCG Outcomes Indicator Set - domain 5				
5.3 Incidence of Healthcare Associated Infection (HCAI) - Methicillin-resistant Staphylococcus aureus (MRSA) (April 2013 - April 2018)	7.57 ●		2.23	19.83
5.4 Incidence of Healthcare Associated Infection (HCAI) - C. difficile (April 2013 - April 2018)	118.0 ●		46	234